arXiv:1209.5289v1 [quant-ph] 24 Sep 2012

Local 3D spin Hamiltonian as a thermally stable surface code

Fabio L. Pedrocchi,[1] Adrian Hutter,[1] James R. Wootton,[1] and Daniel Loss[1]
[1]*Department of Physics, University of Basel, Klingelbergstrasse 82, CH-4056 Basel, Switzerland*

We study a 2D toric code embedded in a 3D Heisenberg ferromagnet in a broken-symmetry state at finite temperature. Stabilizer operators of the toric code are locally coupled to individual spins of the ferromagnet. The effects of the Goldstone modes of the ferromagnet in the ordered phase lead to an energy penalty for anyons that grows linearly with L, the linear size of the toric code. This $O(L)$ energy barrier for logical errors leads to a lifetime of the quantum memory that grows exponentially with L, assuming that the toric code is weakly coupled to a thermal bath with temperature below the phase transition of the ferromagnet. This provides a stable quantum memory with strictly local bounded-strength interactions in less than four dimensions.

PACS numbers: 03.67.Pp,03.67.Lx,05.30.Pr,75.10.Jm

Introduction. Topologically ordered phases of matter like Kitaev's toric code promise the possibility to store and process quantum information in a manner which is resilient against local imperfections [1–3]. However, a finite gap for the creation of topological defects (called *anyons* in the case of the toric code) is not enough to ensure stability against thermal fluctuations [4–7]. If anyons can be created at a constant energy cost and propagate without any further energy penalty, they will at any non-zero temperature destroy the stored quantum information in a time which does not increase with the size of the memory. Indeed, it was shown that not only the toric code but a large class of 1-, 2-, and 3-dimensional Hamiltonians suffer from the aforementioned thermal instability of quantum information [8–10]. This is in contrast to the classical case, where magnetic devices allow the construction of self-correcting hard drives that are stable against both local perturbations and thermal excitations.

Proposals for three-dimensional spin Hamiltonians with local few-spin interactions that do not fall victim to the aforementioned no-go results exist [11–13]. While the energy barrier in Refs. [12] and [13] grows respectively like $O(\log(L))$ [14] and $O(L^{2/3})$, none of these models is expected to allow for a storage time increasing arbitrarily with system size.

Several authors considered the possibility of suppressing the propagation or creation of toric code anyons by coupling them to external fields and found storage times that grow polynomially in L [15–19] or exponentially in L in a certain resonant parameter regime [17, 19]. Suppressing the diffusion of anyons by means of an attractive interaction between them was proposed in Ref. [2] and a possible implementation of such an interaction was found in Ref. [15] by coupling the toric code to a bosonic field. However, the coupling strengths in Ref. [15] diverge polynomially in L. Note that quantum propagation of anyons due to magnetic perturbations in the Hamiltonian can be suppressed as well, namely through the presence of disorder as shown in Refs. [20, 21]. Alternatively, Ref. [16] proposed long-range repulsion between anyons, leading to a vanishing anyon-density for large L. The implementation and benefits of such long-range interactions have been further studied in Refs. [17–19].

In this work, we propose a three-dimensional spin model with purely local interactions of bounded strength that allows for thermally stable storage of quantum information below some critical temperature. We consider a surface code embedded in a bulk ferromagnet and find an anyon chemical potential increasing linearly with L and thus a memory lifetime increasing exponentially with L. This scaling of the lifetime coincides with the four-dimensional toric code [2, 22], which constitutes so far the only known example of a truly self-correcting quantum memory. The origin of this favorable behavior resides in the long-range character of the ferromagnetic transverse susceptibility in the ordered phase since it describes the effective attractive interactions between stabilizer operators. This long-range behaviour is based on the Goldstone modes of the ferromagnet in the broken-symmetry phase. This result is general and does not depend on damping effects and temperature T as long as the ferromagnet stays in the ordered phase. While the long-distance behavior of the susceptibility can be understood by rather general hydrodynamic arguments, we derive an explicit expression for the susceptibility in the one-magnon approximation for the sake of illustration. The effect of the ferromagnet can then be understood in terms of virtual emission and absorption of Goldstone modes (magnons) which mediate long-range interactions between the stabilizer operators. Finally, we show how to choose an appropriate coupling scheme to the ferromagnet in order to hinder hopping of anyons (and not only their creation) by energy barriers that grow proportionally with L.

Model. We introduce two sets of spins, namely $\mathbf{S}_j$ for the spins of the three-dimensional ferromagnet located at site j of a cubic lattice and $\mathbf{I}_j$ for the physical spins-1/2 of the 2D surface code. The Hamiltonian we consider is purely local and takes the following form

$$H = H_{\mathrm{F}} + A \sum_{\mathbf{p}} W_{\mathbf{p}} S^x_{\mathbf{p}} \, , \qquad (1)$$


S-spin
$W_\mathbf{p} = I^z_{\mathbf{p},1} I^y_{\mathbf{p},2} I^z_{\mathbf{p},3} I^y_{\mathbf{p},4}$
2
3
p
1
4
S-spin

FIG. 1. A 2D surface code is embedded in the bulk of a 3D ferromagnet which is in a broken-symmetry state. The stabilizers (plaquettes) $W_\mathbf{p}$ of the surface code locally couple to the S-spins of the ferromagnet (grey dots). Due to the broken symmetry, there exist long-range Goldstone modes or magnons (wiggly line) in the ferromagnet that mediate attractive gravitation-like interactions between the stabilizers.

where A is the coupling constant between the spins of the surface code and the ferromagnet. Here, the plaquette operator $W_\mathbf{p} = I^z_{\mathbf{p},1} I^y_{\mathbf{p},2} I^z_{\mathbf{p},3} I^y_{\mathbf{p},4}$ is the poduct of spins around the square plaquette centered at $\mathbf{p}$ which are defined on a square lattice of linear size L with periodic boundary conditions (we set the lattice constant to unity). The three-dimensional vector $\mathbf{p}$ points towards the spin $\mathbf{S}$ lying in the center of a square plaquette of the surface code, see Fig. 1. Note that this definition of $W_\mathbf{p}$ ensures that the blue and white plaquettes are equivalent to the usual toric code star and plaquette operators [1].

$H_\mathrm{F} = -J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j$ with $J > 0$ is the Hamiltonian of the three-dimensional Heisenberg ferromagnet with the sum restricted to nearest-neighbor sites $\langle i, j \rangle$. The ferromagnet is assumed to be in a broken-symmetry state pointing along the z-direction. In order to implement this formally, an infinitesimal magnetic field in z-direction is added to H_F. The physical spins $\mathbf{I}_j$ of the surface code are embedded in the bulk of the ferromagnet. The interaction Hamiltonian $H_\mathrm{int} = A \sum_\mathbf{p} W_\mathbf{p} S^x_\mathbf{p}$ involves purely local five-body interaction terms. We define the anyon number $n_\mathbf{p}$ such that $W_\mathbf{p} = 1 - 2n_\mathbf{p}$.

For $A \ll J$, we make use of a perturbative Schrieffer-Wolff transformation [23] similar to Ref. [24] to derive the effective plaquette-plaquette interaction (a detailed derivation is given in the Appendix) given by

$$H_\mathrm{eff} = \frac{1}{2} \sum_{\mathbf{p},\mathbf{p}'} J_{\mathbf{p},\mathbf{p}'} W_\mathbf{p} W_{\mathbf{p}'} \ , \tag{2}$$

where the coupling is $J_{\mathbf{p},\mathbf{p}'} = -A^2 \chi_{xx}(\mathbf{p}-\mathbf{p}')$ and $\chi_{\alpha\beta}(\mathbf{r})$ is the static spin susceptibility of the $\mathbf{S}$-spins only. Note that the energy in H_eff can be minimized by either all stabilizers $W_\mathbf{p}$ having a $+1$ or a -1 eigenvalue. Adding the usual toric code Hamiltonian $H_\mathrm{toric} = -\Delta \sum_\mathbf{p} W_\mathbf{p}$ to Hamiltonian (1) would explicitly break this symmetry between anyons and anyon-holes (anyon-holes $\bar{n}_\mathbf{p}$ are defined through $\bar{n}_\mathbf{p} = 1 - n_\mathbf{p}$). However, this term is not needed and its effect is vanishing in the limit of large L, as we discuss in the Appendix, so we neglect it for simplicity. The real space static susceptibility $\chi_{\alpha\beta}(\mathbf{r})$ is defined as the Fourier transform of

$$\chi_{\alpha\beta}(\mathbf{q}, \omega) = i \lim_{\eta \to 0} \int_0^\infty \mathrm{d}t \, e^{(i\omega - \eta)t} \left\langle \left[S^\alpha_\mathbf{q}(t), S^\beta_{-\mathbf{q}} \right] \right\rangle \ , \tag{3}$$

for $\omega = 0$, where $\langle \dots \rangle$ denotes thermal equilibrium expectation values of the $\mathbf{S}$-spins [25] at temperature T. The Fourier components are defined through $S^\alpha_\mathbf{q} = \frac{1}{\sqrt{N_s}} \sum_i e^{-i\mathbf{q}\cdot\mathbf{R}_i} S^\alpha_i$, where N_s is the number of spins in the ferromagnet, and $\mathbf{R}_i$ is a three-dimensional vector denoting the site of spin $\mathbf{S}_i$ of the ferromagnet.

It is not necessary to explicitly calculate the spin susceptibility in the ferromagnetically ordered state to understand its general behavior at large distances (or small $\mathbf{q}$) [26]. Indeed, the spontaneous $SO(3)$ symmetry breaking of the state with finite magnetization pointing along, say, the z-axis, implies the presence of low-frequency Goldstone modes (called magnons in this context) and long-range correlations, i.e., the xx- (and yy-) susceptibility has to diverge for $\mathbf{q} \to \mathbf{0}$ and takes the following generic form in the hydrodynamic regime (low-energy and long wavelength regime) [26]

$$\chi_{xx}(\mathbf{q}, \omega = 0) = \frac{M^2}{R|\mathbf{q}|^2} \text{ for } \mathbf{q} \to \mathbf{0} \ , \tag{4}$$

where $R > 0$ is the stiffness constant of the ferromagnet and $M^2 = |\langle s^z \rangle|^2$ is the magnetization density with $s^z = \frac{1}{N_s} \sum_i S^z_i$. The divergence at $\mathbf{q} \to \mathbf{0}$ is trivially connected with the broken symmetry of the ground state: starting from a ferromagnetic state aligned along the z-direction, the slightest x-magnetic field is able to rotate and align all spins in $x-$direction and thus the response to an external magnetic field indeed diverges at $\mathbf{q} \to \mathbf{0}$. The same line of reasoning applies to the yy-susceptibility. Equation (4) is thus very general and takes into account damping effects due to magnon-magnon interactions [26]. Furthermore, an upper bound for the stiffness constant R can be derived [26],

$$R \leq \frac{1}{3} J \sum_{\langle 0,i \rangle} \langle \mathbf{S}_0 \cdot \mathbf{S}_i - S_0^y S_i^y \rangle \ . \tag{5}$$

Below we give an explicit expression for R in the one-magnon approximation.

The real space static susceptibility now follows by Fourier transforming Eq. (4) which leads to

$$\chi_{xx}(\mathbf{r}) = \frac{M^2}{R} \frac{1}{4\pi |\mathbf{r}|} \ . \tag{6}$$

The plaquette-plaquette interaction is thus long-range gravitation-like and reads

$$J_{\mathbf{p},\mathbf{p}'} = -\frac{A^2 M^2}{4\pi R} \frac{1}{|\mathbf{p} - \mathbf{p}'|} \ . \tag{7}$$

Consequently, Eq. (2) describes a stabilizer Hamiltonian with long-range plaquette-plaquette interactions. Since the stiffness constant R is positive (see also below), the interaction between stabilizer operators $W_{\mathbf{p}}$ is attractive. By rewriting Hamiltonian (2) in terms of anyons we obtain up to some irrelevant constant $(\sum_{\mathbf{p},\mathbf{p}'} J_{\mathbf{p},\mathbf{p}'})$

$$H_{\text{eff}} = -2 \sum_{\mathbf{p},\mathbf{p}'} J_{\mathbf{p},\mathbf{p}'} n_{\mathbf{p}} + 2 \sum_{\mathbf{p},\mathbf{p}'} J_{\mathbf{p},\mathbf{p}'} n_{\mathbf{p}} n_{\mathbf{p}'} \ . \tag{8}$$

Note that for $\mathbf{p} = \mathbf{p}'$ the two summands cancel, such that we can restrict the sums to $\mathbf{p} \neq \mathbf{p}'$. A similar Hamiltonian has been derived in Ref. [17] by coupling Kitaev's honeycomb model to cavity modes. While the non-locality of cavity modes allowed one to obtain constant long-range plaquette-plaquette interactions, here we start from a purely local Hamiltonian and obtain gravitation-like long-range interactions, i.e., $J_{\mathbf{p},\mathbf{p}'} \propto \frac{1}{|\mathbf{p}-\mathbf{p}'|}$. As we discuss below, this is sufficient to stabilize the memory against thermal fluctuations.

For the sake of illustration we calculate the stiffness perturbatively in the one-magnon approximation. For this we make us of the Holstein-Primakoff transformation [27]

$$S_i^z = -S + n_i, \quad S_i^- = \sqrt{2S} \sqrt{1 - \frac{n_i}{2S}} a_i, \ \text{ and } \ S_i^+ = (S_i^-)^\dagger, \tag{9}$$

in the limit $n_i \ll 2S$, where $n_i = a_i^\dagger a_i$. Here, a_i and $a_i^\dagger$ satisfy bosonic commutation relations and the associated quasi-particles are called magnons. In Fourier space, we get $H_F \approx \sum_{\mathbf{q}} \epsilon_{\mathbf{q}} a_{\mathbf{q}}^\dagger a_{\mathbf{q}}$, up to some irrelevant constant, with magnon dispersion $\epsilon_{\mathbf{q}} = 4JS[3 - (\cos(q_x) + \cos(q_y) + \cos(q_z))]$, where we have introduced the Fourier components $a_{\mathbf{q}} = \frac{1}{\sqrt{N}} \sum_i e^{-i\mathbf{q}\cdot\mathbf{R_i}} a_i$. Inserting Eq. (9) into Eq. (3) leads to

$$\begin{aligned} &\chi_{xx}^{(0)}(\mathbf{q}, \omega = 0) = \\ &\lim_{\eta \to 0} \frac{iS}{2} \int_0^\infty \mathrm{d}t\, e^{-\eta t} \left\langle \left[e^{i\epsilon_{\mathbf{q}} t} a_{-\mathbf{q}}^\dagger + e^{-i\epsilon_{\mathbf{q}} t} a_{\mathbf{q}}, a_{\mathbf{q}}^\dagger + a_{-\mathbf{q}} \right] \right\rangle \\ &= \frac{S}{\epsilon_{\mathbf{q}}} \ . \end{aligned} \tag{10}$$

Note that the temperature dependence drops out of the equation above since $\langle [a_{\pm\mathbf{q}}, a_{\pm\mathbf{q}}^\dagger] \rangle = 1$. After a small-$\mathbf{q}$ expansion of $\epsilon_{\mathbf{q}}$ (accurate if one is interested in the physics at large distances), we obtain

$$\chi_{xx}^{(0)}(\mathbf{q}, \omega = 0) = \frac{1}{2J|\mathbf{q}|^2} \ . \tag{11}$$

A comparison between Eqs. (4) and (11) allows us to identify $R^{(0)} = 2JS^2$ since here $(M^{(0)})^2 = S^2$. Note that $R^{(0)}$ saturates the upper bound Eq. (5). We thus obtain that $\chi_{xx}^{(0)}(\mathbf{r}) = \frac{1}{8\pi J |\mathbf{r}|}$, and finally the following approximate plaquette-plaquette coupling

$$J_{\mathbf{p},\mathbf{p}'}^{(0)} = -\frac{A^2 (M^{(0)})^2}{4\pi R^{(0)}} \frac{1}{|\mathbf{p} - \mathbf{p}'|} = -\frac{A^2}{8\pi J |\mathbf{p} - \mathbf{p}'|} \ , \tag{12}$$

which is explicitly attractive since $J > 0$. We point out again that Eq. (12) is the one-magnon approximation of Eq. (7). Notably, the sole effect of both damping due to magnon-magnon interactions and temperature is to renormalize the coefficients of the interaction (12), i.e. $(M^{(0)})^2/R^{(0)} \to M^2/R$, while the form of the gravitational potential is not affected. Indeed, the diverging behavior of the transverse susceptibility at $|\mathbf{q}| \to 0$, see Eq. (4), is a consequence of the broken $SO(3)$-symmetry and is not modified by thermal fluctuations as long as the ferromagnet is in the ordered phase. Besides hydrodynamic and one-magnon approximations, other methods lead to results similar to (4) for the spin susceptibility. In particular, the Tyablikov RPA-approximation [27] is especially useful for a spin-1/2 ferromagnet and leads to the same $\mathbf{q}$-dependence of the susceptibility independent of temperature in the ordered phase. We point out that the creation or annihilation of magnons will not generate anyons in the toric code since the coupling terms in Hamiltonian (1) commute with the toric code stabilizers. The presence of thermal magnons in the ferromagnet will only translate into small fluctuations of the chemical potential for the anyons.

Note that the dimensionality of the ferromagnet used here is a central ingredient since Heisenberg ferromagnets in lower dimensions do not undergo a phase transition at $T > 0$ [28].

Thermally Stable Quantum Memory. In this section we show that the long-range attraction (7) between $W_{\mathbf{p}}$ operators leads to a stable quantum memory. Indeed, inserting Eq. (7) into Hamiltonian (8) and using the fact that in the continuum limit $\sum_{\mathbf{p}\neq\mathbf{0}} \frac{1}{|\mathbf{p}|} \approx \int_{L\times L} \frac{1}{|\mathbf{p}|}\, \mathrm{d}\mathbf{p} =$

cL, with a geometrical factor $c \in O(1)$, we obtain

$$H = \mu(L) \sum_{\mathbf{p}} n_{\mathbf{p}} + 2 \sum_{\mathbf{p} \neq \mathbf{p}'} J_{\mathbf{p},\mathbf{p}'} n_{\mathbf{p}} n_{\mathbf{p}'} \ , \tag{13}$$

where the chemical potential of the anyons is

$$\mu(L) = \frac{cA^2 M^2}{2\pi R} \cdot L \tag{14}$$

and thus increases linearly with the system size L. This implies a quantum information storage time that grows exponentially in L and β, where $\beta = 1/T$ is the inverse temperature of a bath weakly coupled to the memory, as originally demonstrated in Ref. [29] assuming that the interaction with the thermal bath can be described by the Davies equation. Below we present alternative arguments leading to the same conclusion.

The second term in Eq. (13) describes a gravitation-like attraction between anyons, which is much weaker than the logarithmic potential found in Ref. [15] (indeed, the energy needed for the anyons to escape the gravitational potential does not increase with L). Since this term helps to keep newly created anyon pairs attached to each other (for temperatures below the interaction strength $A^2M^2/\pi R$), it will have a further beneficial effect on the memory lifetime. Note that this second term effectively reduces the anyon chemical potential. However, this reduction is negligible since the anyon density is exponentially suppressed by the first term.

Let us try to understand in more detail the decoherence process of the memory in contact with a simple model of a bath. We assume that the bath supports single-spin processes in which an energy ω is transfered from the anyon system to the bath with rate $\gamma(\omega)$ and that $\gamma(0) \neq 0$ [30]. Let $\delta(N)$ denote the average cost to create an anyon pair if there are already N pairs present. The gravitational interaction will lead to $\delta(N \geq 1) < \delta(0) = 2\mu(L) - A^2M^2/(\pi R)$. However, in the Appendix we show that this reduction will not lead to a finite self-consistent number of anyon pairs and that in fact we will have $\delta(N \geq 1) \approx \delta(0)$ in the relevant regime.

Since the presence of only two anyons diffusing across the memory leads to an uncorrectable logical error in times of order $L^2/\gamma(0)$ [16], we need to show that the time for the creation of two nearby anyons that are not directly annihilated increases exponentially with system size. Whenever a new pair of anyons is created, their total hopping rate is given by $6\gamma(0)$ [31] such that the probability that one of the two anyons ever moves before the pair gets annihilated is $6\gamma(0)/[\gamma(\delta(0)) + 6\gamma(0)]$. Since $\gamma(\delta(0)) = \exp(\beta\delta(0))\gamma(-\delta(0))$ (which follows from the detailed balance condition) and the code consists of L^2 physical spins, we conclude that the total rate for creation of anyon pairs that do not directly get annihilated is given by

$$L^2 \gamma(-\delta(0)) \frac{6\gamma(0)}{\gamma(\delta(0)) + 6\gamma(0)} \leq 6L^2 e^{-\beta\delta(0)} \gamma(0) \ . \tag{15}$$

The time needed to create such a pair is thus of order $\exp(\beta\delta(0))/L^2\gamma(0)$. In conclusion, we found a lower bound for the quantum memory storage time that increases exponentially with system size.

This lifetime does not apply if the initial state of the system has anyons already present. Suppose that errors occur during preparation of the initial state, creating a finite density of anyons. If these errors are sufficiently sparse, it will be possible for error correction to recover the initial state. It is the job of the Hamiltonian to preserve this error correctability until the desired time of readout. The Hamiltonian given above will energetically favour the annihilation of anyons on neighboring plaquettes, undoing some of the errors. However, we can expect that a finite density of pairs will have been non-neighbouring, and so will remain. These only need to diffuse a constant distance to make correction ambiguous, which leads to a constant lifetime for the memory. To prevent this we can split the plaquettes into two types. 'Strongly coupled' plaquettes are coupled to the ferromagnet with a strength A_s. 'Weakly coupled' plaquettes have a strength $A_w < A_s$. These are chosen such that any sequence of single- or local two-spin errors that move an anyon from one weakly coupled plaquette to another must move it via a strongly coupled plaquette (example patterns are given in the Appendix). The chemical potential for the plaquettes will change from the form in Eq. (14), giving different values $\mu_s(L)$ and $\mu_w(L)$ for the two types of plaquette. Performing the summation (as described in the Appendix) shows that the factor A^2 becomes $A_s\bar{A}$ for $\mu_s(L)$ and $A_w\bar{A}$ for $\mu_w(L)$ ($\bar{A}$ being a weighted average). The energy barrier required for anyon movement is therefore of order $(1 - A_w/A_s)\mu_s(L)$, which increases linearly with system size. The resulting suppression of diffusion leads to a lifetime that increases exponentially with system size, even when the initial state has a finite density of anyons.

It may come as a surprise that associating some stabilizers with a lower energy penalty has a beneficial effect on the memory. However, note that the weakly coupled plaquettes allow energy to be dissipated from the anyons to the bath by hopping of an anyon from a strongly to a weakly coupled plaquette. On the other hand, if the chemical potential is independent of the anyon position, as in Eq. (14), this is only possible through annihilation of anyons.

Conclusions. To the best of our knowledge, we have presented the first Hamiltonian in less than four dimensions that involves only local few-spin interactions of bounded strength and leads to a stable quantum memory. We showed that the the presence of a ferromagnet stabilizes the memory by increasing the anyon's chemical potential linearly with the system size, and thus leads to a lifetime of the memory that grows exponentially with L. An anyon chemical potential that is proportional to L trivially implies a corresponding energy gap between different groundstates of the toric code. This is not in contradiction to no-go results [8–10] since our starting Hamiltonian (1) is not a stabilizer Hamiltonian and the effective stabilizer Hamiltonian (2) is highly non-local.

The question of how to engineer the five-body interactions of Hamiltonian (1) remains open. One possibility may be to couple the spins of the ferromagnet to Kitaev's honeycomb model [32] (since it is equivalent to the toric code up to fourth order in perturbation theory), similar to Ref. [17]. To conclude we would like to point out that our analysis is applicable to any stabilizer Hamiltonian and not restricted to the toric code. Indeed, the sole property which is needed to derive our results is that all stabilizer operators $W_{\mathbf{p}}$ commute among themselves and get coupled by massless Goldstone modes.

Acknowledgements. We would like to thank D. DiVincenzo, F. Hassler, B. Terhal, and L. Trifunovic for helpful discussions. This work was supported by the Swiss NSF, NCCR Nanoscience, and NCCR QSIT.

---

[1] A. Kitaev, Ann. Phys. **303**, 230, (2003).
[2] E. Dennis, A. Kitaev, A. Landahl, and J. Preskill, J. Math. Phys. **43**, 44524505 (2002).
[3] S. Bravyi, M. Hastings, and S. Michalakis, J. Math. Phys. **51** 093512 (2010).
[4] Z. Nussinov and G. Ortiz, Phys. Rev. B **77**, 064302 (2008).
[5] C. Castelnovo and C. Chamon, Phys. Rev. B **76**, 184442 (2007).
[6] R. Alicki, M. Fannes, and M. Horodecki, J. Phys. A: Math. Theor. **40**, 6451 (2007).
[7] R. Alicki, M. Fannes, and M. Horodecki, J. Phys. A: Math. Theor. **42**, 065303 (2009).
[8] S. Bravyi and B. Terhal, New J. Phys. **11**, 043029 (2009).
[9] J. Haah and J. Preskill, arXiv:1011.3529 (2010).
[10] B. Yoshida, Ann. Phys. **326**, 2566 (2011).
[11] D. Bacon, Phys. Rev. A **73**, 012340 (2006).
[12] J. Haah, Phys. Rev. A **83**, 042330 (2011).
[13] K. Michnicki, arXiv:1208.3496 (2012).
[14] S. Bravyi and J. Haah, Phys. Rev. Lett. **107**, 150504 (2011).
[15] A. Hamma, C. Castelnovo, and C. Chamon, Phys. Rev. B **79**, 245122 (2009).
[16] S. Chesi, B. Röthlisberger, and D. Loss, Phys. Rev. A **82**, 022305 (2010).
[17] F. L. Pedrocchi, S. Chesi, and D. Loss, Phys. Rev. B **83**, 115415 (2011).
[18] B. Röthlisberger, J. R. Wootton, R. M. Heath, J. K. Pachos, and D. Loss, Phys. Rev. A **85**, 022313 (2012).
[19] A. Hutter, J. R. Wootton, B. Röthlisberger, and D. Loss, arXiv:1206.0991 (2012).
[20] J. R. Wootton and J. K. Pachos, Phys. Rev. Lett. **107**, 030503 (2011).
[21] C. Stark, L. Pollet, A. Imamoglu, and R. Renner, Phys. Rev. Lett. **107**, 030504 (2011).
[22] R. Alicki, M. Horodecki, P. Horodecki, and R. Horodecki, Open Syst. Inf. Dyn. **17**, 1 (2010).
[23] S. Bravyi, David DiVincenzo, and D. Loss, Ann. Phys. **326**, 2793 (2011).
[24] P. Simon, B. Braunecker, and D. Loss, Phys. Rev. B **77**, 045108 (2008).
[25] Since we calculate $H_{\text{eff}} \propto A^2/J$ to lowest order only, we can omit any A-dependence in $\langle\ldots\rangle$. This amounts to neglect backaction effects (from the surface code on the ferromagnet), which is justified for $A/J \ll 1$. Additional backaction effects would result from higher order terms in the Schrieffer-Wolff transformation. Moreover, backaction effects from a 2D surface on a bulk system become negligible in the thermodynamic limit.
[26] D. Forster, *Hydrodynamic Fluctuations, Broken Symmetry, and Correlation Functions* (Benjamin, MA 1975).
[27] W. Nolting and A. Ramakanth, *Quantum Theory of Magnetism* (Springer, Berlin, 2009).
[28] N.D. Mermin and H. Wagner, Phys. Rev. Lett. **17**, 1133 (1966).
[29] S. Chesi, D. Loss, S. Bravyi, and B. M. Terhal, New J. Phys. **12**, 025013 (2010).
[30] If we had $\gamma(0) = 0$, as is the case for super-Ohmic baths, this would of course have a greatly beneficial influence on the memory lifetime as it forbids direct hopping processes of anyons. See [16] for more details about the decoherence of quantum memories under the influence of super-Ohmic baths.
[31] Strictly speaking, the energy cost for hopping is greater than zero since it increases the potential energy in the gravitational potential. However, we approximate this energy cost by zero for simplicity, neglecting the beneficial effect of the anyon attraction and obtaining a lower bound on the actual lifetime.
[32] A. Kitaev, Ann. Phys. **321**, 2 (2006).

APPENDIX

Interactions mediated by a translationally invariant system. For the sake of completeness, we show here a detailed derivation of Eq. (2) of the main text with the use of a perturbative Schrieffer-Wolff transformation similar to Ref. [1]. Let us thus consider the Hamiltonian

$$H = H_0 + V = H_0 + \sum_{\mathbf{q}} S^{\alpha}_{\mathbf{q}} A_{-\mathbf{q}} \,, \tag{16}$$

where H_0 is a general $\mathbf{S}$-spin Hamiltonian and A_i arbitrary operators which commute with H_0 and with each other. The Fourier components are defined through $S_{\mathbf{q}} = \frac{1}{\sqrt{N_s}} \sum_i e^{-i\mathbf{q}\cdot\mathbf{R}_i} \mathbf{S}_i$ and $A_{\mathbf{q}} = \frac{1}{\sqrt{N_s}} \sum_i e^{-\mathbf{q}\cdot\mathbf{R}_i} A_i$, where N_s denotes the number of spins $\mathbf{S}_i$ and $\mathbf{R}_i$ their site. The effective Hamiltonian up to second order in A is then simply given by

$$H^{(2)}_{\text{eff}} = H_0 + U = H_0 - \frac{i}{2} \lim_{\eta \to 0} \int_0^{\infty} \mathrm{d}t \, e^{-\eta t} \left[V(t), V\right] \,, \tag{17}$$

where $V(t) = e^{iH_0 t} V e^{-iH_0 t}$ is given in the Heisenberg representation ($\hbar = 1$). If we now insert the explicit expression for V we obtain

$$\begin{aligned} U &= -\frac{i}{2}\lim_{\eta\to 0}\sum_{\mathbf{q},\mathbf{q}'}\int_0^\infty \mathrm{d}t\, e^{-\eta t}\left[S^\alpha_{\mathbf{q}}(t)A_{-\mathbf{q}}, S^\alpha_{\mathbf{q}'}A_{-\mathbf{q}'}\right] \\ &= -\frac{i}{2}\lim_{\eta\to 0}\sum_{\mathbf{q},\mathbf{q}'}\int_0^\infty \mathrm{d}t\, e^{-\eta t}([S^\alpha_{\mathbf{q}}(t), S^\alpha_{\mathbf{q}'}]A_{-\mathbf{q}'}A_{-\mathbf{q}} \\ &\qquad + S^\alpha_{\mathbf{q}}(t)S^\alpha_{\mathbf{q}'}\underbrace{[A_{-\mathbf{q}}, A_{-\mathbf{q}'}]}_{=0}) \; . \end{aligned} \tag{18}$$

We assume that the $\mathbf{S}$-spins are in thermal equilibrium, described by the canonical density matrix $\rho = e^{-\beta H_F}/\mathrm{Tr}\, e^{-\beta H_F}$, where H_F is the $\mathbf{S}$-spin Hamiltonian without the coupling to the plaquettes and corresponds to the main part of the Hamiltonian in Eq. (16), i.e., $H_F = H_0$. In doing so, we neglect the backaction of the surface code on the ferromagnet. This is justified for two reasons. First, to include the coupling term $\propto A$ in ρ would lead to higher order corrections in the coupling constant A (we retain only the leading order terms up to order A^2 in the effective Hamiltonian H_{eff}). Second, the surface code, being a two-dimensional lattice, acts only as a boundary perturbation on the three-dimensional ferromagnet, and thus this perturbation becomes negligibly small in the thermodynamic limit. Further, formally we need to add an infinitesimal fictitious Zeeman field to H_F that breaks the $SO(3)$ symmetry explicitly and induces a spontaneous ferromagnetic order below a critical temperature. Such a field also regularizes the zero-mode of the ferromagnet and thus makes the Schrieffer-Wolff series well-defined formally. After taking the thermodynamic limit, we then let this field go to zero in expectation values of physical quantities. Here, we rely on a formal perturbation expansion in powers of A/J. Convergence of this formal expansion is an interesting question by itself and can be approached along the lines lines discussed in Ref. [2]. However, such rigorous treatment is beyond the present scope and will be addressed elsewhere.

The equilibrium expectation values are then denoted by $\langle\ldots\rangle$. If H_0 is translationally invariant, such that $\langle S^\alpha_{\mathbf{r}_i}S^\beta_{\mathbf{r}_j}\rangle = \langle S^\alpha_{\mathbf{0}}S^\alpha_{\mathbf{r}_j-\mathbf{r}_i}\rangle$, we have $\langle S^\alpha_{\mathbf{q}}S^\alpha_{\mathbf{q}'}\rangle = \langle S^\alpha_{\mathbf{q}}S^\alpha_{-\mathbf{q}}\rangle\delta_{\mathbf{q}+\mathbf{q}',\mathbf{0}}$, and thus

$$\begin{aligned} U &= -\frac{i}{2}\lim_{\eta\to 0}\sum_{\mathbf{q}}\int_0^\infty \mathrm{d}t\, e^{-\eta t}\left\langle[S^\alpha_{\mathbf{q}}(t), S^\alpha_{-\mathbf{q}}]\right\rangle A_{\mathbf{q}}A_{-\mathbf{q}} \\ &= -\frac{1}{2}\sum_{\mathbf{q}} A_{-\mathbf{q}}\chi_{\alpha\alpha}(\mathbf{q})A_{\mathbf{q}} \; , \end{aligned} \tag{19}$$

where $\chi_{\alpha\alpha}(\mathbf{q})$ is the static spin susceptibility.

Mean-field analysis of anyon pair creation. Assume that there are already N anyon *pairs* present. We want to determine the average (averaged over all possible positions of the existing anyons) energy cost $\delta(N)$ to create a new pair. From the point of view of one of the two newly created anyons, we assume that the existing $2N$ anyons are uniformly distributed over all L^2-2 remaining positions. The averaged interaction between one of the newly created anyons and each existing one is thus

$$\begin{aligned} &\frac{1}{L^2-2}\left(2\cdot 2\sum_{\mathbf{p}\neq\mathbf{0}} J_{\mathbf{p},\mathbf{0}} + A^2M^2/(\pi R)\right) \\ &\quad = -\frac{1}{L^2-2}\left(2\mu(L) - A^2M^2/(\pi R)\right) \; , \end{aligned} \tag{20}$$

where we have subtract the energy $-A^2M^2/(\pi R)$ due to attraction with the other anyon of the same pair. Indeed, we are only intersted in the attraction energy due to anyons which are already present before the creation of the pair. The total energy $\delta(N)$ to create the new pair is thus given by

$$\begin{aligned} \delta(N) &= \delta(0) - \frac{4N}{L^2-2}\left(2\mu(L) - A^2M^2/(\pi R)\right) \\ &= \delta(0)\left(1 - \frac{4N}{L^2-2}\right) \; , \end{aligned} \tag{21}$$

where $\delta(0) = 2\mu(L) - A^2M^2/(\pi R)$.

The mean-field energy of N anyon pairs is thus

$$\begin{aligned} E_{\mathrm{mf}}(N) &= \sum_{i=0}^{N-1}\delta(i) \\ &= \delta(0)N\frac{L^2-2N}{L^2-2} \; . \end{aligned} \tag{22}$$

The symmetry $N \leftrightarrow L^2/2 - N$ is reminiscent of the fact that the energy in Eq. (2) of the main text can be minimized by either all stabilizers having a $+1$ eigenvalue (no anyons present) or a -1 eigenvalue (memory full of anyons). The energetic gap between the sector in which there are almost no anyons and the sector in which the memory is full of anyons is of order $\Delta(0)L^2 = O(L^3)$, so transitions between these two sectors happen on timescales much longer than the time before the stored quantum information is lost. Consequently, each sector may serve as a thermally stable quantum memory, but at each moment in time we can only use one of the two. Without loss of generality, we consider the case where the sector with (almost) no anyons present is used for quantum information storage.

From Eq. (21) we have that $\delta(N) = \delta(0)(1-2n)$, where n denotes the density of anyons. As there can only be zero or one anyon at each position, we obtain the self-consistent equation for the mean-field anyon density *in equilibrium*

$$n_{\mathrm{mf}} = \left[\exp\left(\beta\delta(0)(1-2n_{\mathrm{mf}})\right) + 1\right]^{-1} \; . \tag{23}$$

If the left-hand side of this equation is smaller/larger than the right-hand side, the anyon density will tend to increase/decrease. If n_{mf} solves this equation, so does

$1-n_{\mathrm{mf}}$. One self-consistent density is $n_{\mathrm{mf}}=\frac{1}{2}$. The stability of this density depends on the temperature of the bath. For $\beta\delta(0)<2$ we have a unique self-consistent density $n_{\mathrm{mf}}=\frac{1}{2}$ and this density is also stable. For $\beta\delta(0)>2$ the density $\frac{1}{2}$ becomes unstable and two new stable self-consistent densities n^* and $1-n^*$ emerge (let n^* denote the smaller of the two). The system of anyons therefore shows a phase transition and spontaneous breaking of the anyon anyon-hole symmetry at a critical temperature $\delta(0)/2$, which is of order $\frac{A^2}{J}L$. However, this temperature becomes only relevant if it is lower than the critical temperature of the ferromagnet which is of order J, which will not be the case in the limit of large L. For the purpose of quantum information storage, we are clearly interested in temperatures below both of these critical temperatures.

Adding the toric code Hamiltonian $H_{\mathrm{toric}} = -\Delta\sum_{\mathbf{p}} W_{\mathbf{p}}$ to Eq. (1) in the main text explicitly breaks the symmetry between anyons and anyon holes and will lead to an additional summand $4N\Delta$ in Eq. (22). However, the modification of the self-consistent densities n^*, $1-n^*$, and $\frac{1}{2}$ through this new term becomes vanishing for large L, as Δ does unlike $\delta(0)$ not grow with L.

Let us consider the self-consistent solution n^*. We want to show that n^* is exponentially suppressed with L and consequently that the number of anyons itself goes to zero in the thermodynamic limit. After straightforward algebra, one can show that $n=2e^{-\beta\delta(0)}<1/2$ with $\beta\delta(0)e^{-\beta\delta(0)}<\frac{\log(2)}{4}$ (note that this condition is readily satisfied since $\delta(0)$ grows linearly with L) satisfies

$$[\exp(\beta\Delta(0)(1-2n))+1]^{-1} < n \;, \tag{24}$$

and therefore $n>n^*$. Since n is by definition exponentially suppressed with L and $n^*<n$ we finally conclude that the self-consistent solution n^* of Eq. (23) goes exponentially to zero with L. A direct consequence of this is that the equilibrium number of anyons n^*L^2 also vanishes exponentially with L and will generally be much smaller than the minimal positive value 2. Hence the anyon number will fluctuate between 0 and small even integers, such that $\delta(N)\approx\delta(0)$ from Eq. (21).

An example pattern for strongly and weakly coupled plaquettes. In the toric code model there are two types of anyon, e and m, which reside on two kinds of plaquette, s and p, respectively. Note that, when the code is defined with spins on the edges of the lattices, these correspond to the stars and plaquettes, respectively.

Consider a spin in the square tiling of Fig. 2 (a), shared by two s-plaquettes to the top-left and bottom-right and two p-plaquettes to the top-right and bottom-left. The application of a Pauli I^z to such a spin will affect the e anyon occupations of the two s-plaquettes. If both were initially empty, an anyon pair will be created. If both initially held an anyon, this pair will be annihilated. If only one held an anyon, it will be moved to the other plaquette. The application of a Pauli I^y has the same effect for the m anyons of the p-plaquettes. For spins where the positions of s- and p-plaquettes are exchanged, the roles of I^z and I^y are also exchanged. No operation exists that can move an anyon from an s-plaquette to a p-plaquette, or vice-versa.



FIG. 2. Two tilings of plaquettes are shown on which the code may be defined. Spins are located on vertices. (a) The square tiling, as used in the main text. s-plaquettes are shown in dark blue, blue, or light blue, p-plaquettes are shown in white. (b) An alternative tiling, with alternating triangular and octagonal plaquettes. s-plaquettes are shown in dark blue and blue, p-plaquettes are shown in white and grey.

Creation, movement and annilation of anyons are therefore achieved by Pauli operations. Using single spin operations, creation of a pair will always lead to the anyons occupying neighboring plaquettes (where neighoring means that they share exactly one spin). Similarly, single spin operations can only move anyons from one plaquette to a neigboring one, or annihilate anyons on neighboring plaquettes. Since we assume that the system-bath coupling supports only single spin errors, it is exactly these processes that we consider during thermalization. However, it should be remembered that two-spin perturbations may also be present in the Hamiltonian. Local two-spin errors should also be considered, which can create, annihilate and transport anyons on next-to-neighboring plaquettes.

With this in mind, we wish to split both s- and p-plaquettes into two groups, one of which will be strongly coupled to the ferromagnet with a coupling A_s and the other of which will be weakly coupled with a strength $A_w<A_s$. This will give the plaquettes of the former a higher chemical potential than those of the latter, with an energy difference that increases linearly with system size.

The pattern of strongly and weakly coupled plaquettes should be chosen such that anyons become trapped within the latter, which will occur if two conditions are satisfied. Firstly, any anyons initially on strongly coupled plaquettes should quickly move into a nearby weakly coupled plaquette. Secondly, it should not be possible for anyons to be moved from one weakly coupled plaquette (or a small cluster of weakly coupled plaquettes) to another by a sequence of either single- or two-spin operations without passing through a strongly coupled plaquette.

The first condition can be met if anyons on strongly coupled plaquettes cannot be moved large distances by a sequence of either single or two spin operations without either moving through a weakly coupled plaquette, or through a strongly coupled plaquette that neigbors a weakly coupled one. The latter is relevant because the difference in chemical potential between the two types of plaquette will exponentially suppress the probability that the anyon will do anything but decay into the weakly coupled plaquette in this case.

Both conditions are satisfied by the pattern shown in Fig. 2 (a). Here, weakly coupled s-plaquettes are shown in dark blue. Strongly coupled s-plaquettes that neighbor weakly coupled s-plaquettes are shown in blue, and those that do not are shown in light blue. Regions of strongly coupled plaquettes that do not neighbor weakly coupled plaquettes are separated from each other by a width of three spins. Sequences of one- and two-spin operations therefore cannot move anyons in one such region to another without going via strongly coupled plaquettes that do neighbor weakly coupled plaquettes, which will almost certainly result in the anyon decaying into the neighboring weakly coupled plaquettes. Similarly, regions of weakly coupled plaquettes are separated by the same width, preventing movement between them without going via strongly coupled plaquettes.

The initial movement of anyons on strongly coupled plaquettes to nearby weakly coupled plaquettes may cause ambiguity for error correction if the error rate during initialization is too high. Even so, for sufficiently low error rates this movement will have no effect on correctability. Once the movement is complete, the exponential suppression of diffusion will then ensure that the correctability of the errors is preserved for a time exponential with the system size, since such an exponentially long timescale will be required for the anyons to climb out of the weakly coupled plaquettes.

We will now demonstrate that the difference in chemical potentials between strongly and weakly coupled plaquettes leads to the energy barrier required to suppress diffusion. To determine the chemical potential of an arbitrary plaquette $\mathbf{p}$ (which is either s- or p-type), the following sum over all plaquettes must be performed

$$\mu_{\mathbf{p}}(L) = \frac{M^2}{2\pi R} A_{\mathbf{p}} \sum_{\mathbf{p}'}{}' A_{\mathbf{p}'} \frac{1}{|\mathbf{p}-\mathbf{p}'|} \ , \qquad (25)$$

where the prime in $\sum'$ means that $\mathbf{p}' \neq \mathbf{p}$. Here $A_{\mathbf{p}'}$ denotes the coupling of plaquette $\mathbf{p}'$ which will be A_s or A_w depending on whether this plaquette is weakly or strongly coupled, respectively. By numerically performing the summation we find that, in the $L \to \infty$ limit, it takes the form

$$\sum_{\mathbf{p}'}{}' A_{\mathbf{p}'} \frac{1}{|\mathbf{p}-\mathbf{p}'|} = \frac{3A_s + A_w}{4} c \cdot L \ , \qquad (26)$$

where c is an $O(1)$ constant, as mentioned in the main text. Its value does not depend on whether the sum is centered on a strongly or weakly coupled plaquette, or at least does not do so to a non-negligible degree. To three decimal places its value is found numerically to be $c = 3.524$. The linear combination of A_s and A_w is a weighted average $\bar{A} = (3A_s + A_w)/4$, which arises from the fact that there are three times as many strongly coupled plaquettes as weakly coupled plaquettes. The chemical potentials for weakly and strongly coupled plaquettes are then

$$\mu_s(L) = \frac{cA_s\bar{A}M^2}{2\pi R} \cdot L \ , \quad \mu_w(L) = \frac{cA_w\bar{A}M^2}{2\pi R} \cdot L \ . \quad (27)$$

Clearly, $\mu_s(L) - \mu_w(L) = O(L)$, giving the required energy barrier.

Alternative tiling with four-body coupling. A pattern of strongly and weakly coupled plaquettes, stable against single-spin errors, is shown for an alternative tiling in Fig. 2 (b). Strongly (weakly) coupled s-plaquettes are shown in blue (dark blue) and strongly (weakly) coupled p-plaquettes are shown in white (grey). For this tiling it is still true that e anyons can only be created and moved between neighboring s-plaquettes, and m anyons between neighboring p-plaquettes. Note that all strongly coupled plaquettes in this tiling are triangular. The $W_{\mathbf{p}}$ for these will therefore be three-body operators, making the code-ferromagnet coupling only a four-body term. On the other hand, weakly coupled plaquettes are octagons with eight-body $W_{\mathbf{p}}$ and nine-body terms required for the code-ferromagnet coupling. Since these many-body terms will most likely be generated by perturbative methods, with a higher number of spins in a term generated by higher orders of perturbation theory, the difference in coupling strengths will arise naturally.

Due to the practical difficultly in generating many-body terms, we can consider not coupling the octagonal plaquettes to the ferromagnet. Only the four-body terms required to coupled the triangles are then needed, which should be easier to implement than the five-body terms required for the square tiling. Despite the fact that only a fraction of the plaquettes are coupled to the ferromagnet, the memory is still stable against thermal errors. This is because any single spin error must still create at least one anyon on, or move anyons through, energetically penalized triangular plaquettes. The energy barrier that increases linearly with system size is therefore still intact, and ensures that anyon creation and diffusion are exponentially suppressed.

Unfortunately, stability against local Hamiltonian perturbations does not remain strong without the coupling of octagons. Without an energy penalty, two-body perturbations are free to create and move anyons between next-to-neighboring octagonal plaquettes. This avoids the energy barrier and so leads to uncorrectable errors in a constant time. However, it is possible to avoid this by carefully considering what types of perturbation are present, and then designing the $W_{\mathbf{p}}$ such that they are unable to perform such hopping processes. For example, let us use $W_{\mathbf{p}} = I^x_{\mathbf{p},1} I^y_{\mathbf{p},2} I^z_{\mathbf{p},3}$ for triangular s-plaquettes. Here spin 1 is that shared with the neighboring triangular s-plaquette and the numbering proceeds clockwise. Let us also use $W_{\mathbf{p}} = I^z_{\mathbf{p},1} I^y_{\mathbf{p},2} I^x_{\mathbf{p},3}$ for triangular p-plaquettes with corresponding numbering. No nearest neighbor isotropic perturbation of the form $I^\alpha_i I^\alpha_j$, for $\alpha \in \{x, y, z\}$, commutes with all of these operators. This means such perturbations will be suppressed by the energy barrier and will not be able to move anyons between octagonal plaquettes. If only perturbations of this form are present in the system, the memory will remain stable.

[1] P. Simon, B. Braunecker, and D. Loss, Phys. Rev. B **77**, 045108 (2008).

[2] S. Bravyi, David DiVincenzo, and D. Loss, Ann. Phys. **326**, 2793 (2011).